

REPORT DOCUMENTATION PAGE*Form Approved*
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 19-22, 1995	3. REPORT TYPE AND DATES COVERED Annual Symposium, 1995	
4. TITLE AND SUBTITLE Degradation of Munitions and Chlorinated Solvents by Aquatic Plants			5. FUNDING NUMBERS N/A	
6. AUTHOR(S) N. Lee Wolfe				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Environmental Research Laboratory U.S. EPA, 960 College Station Road, Athens, GA 30605			8. PERFORMING ORGANIZATION REPORT NUMBER N/A	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) SERDP 901 North Stuart St. Suite 303 Arlington, VA 22203			10. SPONSORING / MONITORING AGENCY REPORT NUMBER N/A	
11. SUPPLEMENTARY NOTES Presented at Fourteenth Annual Symposium 1995, Current Topics in Plant Biochemistry, Physiology and Molecular Biology, <i>Will Plants Have a Role in Bioremediation?</i> , University of Missouri, Columbia, MO, pp. 45-46, April 19-22, 1995. No copyright is asserted in the United States under Title 17, U.S. code. The U.S. Government has a royalty-free license to exercise all rights under the copyright claimed herein for Government purposes. All other rights are reserved by the copyright owner.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited			12b. DISTRIBUTION CODE A	
13. ABSTRACT (Maximum 200 Words) Nitroreductase and dehalogenase enzymes have been isolated from sediments and soils. Using enzyme linked immunospecific assays (ELISA), a number of aquatic plants have been identified as sources of the enzymes. The plants were then brought back into the laboratory and evaluated as candidates for further remediation studies.				
14. SUBJECT TERMS nitroreductase, dehalogenase, ELISA, aquatic plants, SERDP			15. NUMBER OF PAGES 2	
			16. PRICE CODE N/A	
17. SECURITY CLASSIFICATION OF REPORT unclass	18. SECURITY CLASSIFICATION OF THIS PAGE unclass	19. SECURITY CLASSIFICATION OF ABSTRACT unclass	20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Fourteenth Annual Symposium

1995

*Current Topics in Plant Biochemistry,
Physiology and Molecular Biology*

19980806 056

Will Plants Have a Role in Bioremediation?



Heavy Metals: Accumulators, Tolerance and Exclusion
Mechanisms

◆
Plant Assisted Decontamination

◆
Plant Assisted Degradation of Organics

◆
Ecological Considerations of Plant-Based Remediation

Abstract Book



FOURTEENTH ANNUAL SYMPOSIUM

**Current Topics in Plant Biochemistry, Physiology and Molecular
Biology**

April 19-22, 1995

**VIDEO TAPING OF LECTURES IS
FORBIDDEN WITHOUT PERMISSION OF
EACH SPEAKER.**

**PHOTOGRAPHY OF POSTERS IS
FORBIDDEN WITHOUT PERMISSION OF
PRESENTERS.**

Degradation of Munitions and Chlorinated Solvents by Aquatic Plants

N. Lee Wolfe

Environmental Research Laboratory
U. S. Environmental Protection Agency
960 College Station Road
Athens, GA 30605

Nitroreductase and dehalogenase enzymes have been isolated from sediments and soils. Using enzyme linked immunospecific assays (ELISA), a number of aquatic plants have been identified as sources of the enzymes. The plants were then brought back into the laboratory and evaluated as candidates for further remediation studies.

In the case of the nitroreductase, detailed studies with Parrot Feather, Stonewort, Hornwort and Spyrigia in water and sediment systems have shown that the plants readily degrade munitions. Significant findings are:

- (1) The pathway for TNT degradation with a plant nitroreductase enzyme has been described.
- (2) A variety of aquatic and terrestrial plants containing the nitroreductase enzyme have been shown to degrade TNT.
- (3) RDX and HMX also undergo degradation.
- (4) Other nitroaromatic compounds associated with TNT manufacturing are also degraded by the plant enzyme system.
- (5) Monoaminodinitro, diaminomononitro substituted toluenes and benzenes are readily degraded by the enzyme system.
- (6) No amino coupling products have been detected in any of the systems studied to date, i.e., azo or azoxy compounds.
- (7) The kinetics of the degradation in water are very fast with disappearance half-lives on the order of minutes for TNT.
- (8) Mass balances and reaction pathways have been described and the "final fate" of ^{14}C ring labeled TNT has been described.
- (9) ^{14}C labeled ring opened products are incorporated in the plant material.
- (10) Aquatic plants with high nitroreductase and laccase activity have been shown to grow well in soil/water systems using soils with high concentrations of TNT (5000 ppm in the soil).
- (11) The nitro reductase enzyme system is stable in sediments for months in both oxic and anoxic sediments. The enzyme can be isolated from sediments with high organic carbon

with the sediments having the highest organic carbon having the highest enzyme activity. (12) After equilibration of soil contaminated TNT with water and plants no TNT or related compounds could be detected in the aqueous phase even after 8 weeks. (13) The plants are effective in degrading TNT, RDX and HMX in water to below 1 ppb. (14) Using an ELISA field test, indigenous plants at a given site can be identified that degrade TNT.

Based on scaled up bench-top results, the process has been further studied in the field in scaled up batch systems. The results have been consistent with the laboratory findings. In the field study, using TNT contaminated sediment/water/Parrot Feather systems, it was possible to drive the aqueous phase TNT and the transient amino intermediates to below 0.1 ppm (the limit of detection in these studies). In separate studies, parrot feather was shown to reduce TNT and the corresponding intermediates to concentrations below 1 ppb.

Similarly for the dehalogenase enzyme, laboratory studies with *Spyrigia* and Parrot Feather have shown that the plants can degrade a number of chlorinated aliphatic compounds including the chlorinated solvents TCE and PCE. The pathways, mass balances and kinetics of the reactions have been elucidated.